

## MILLIMETER WAVE SPECTROSCOPY OF RYDBERG STATES OF MOLECULES IN THE REGION OF 260-295 GHz

DAVID GRIMES, *Department of Chemistry, MIT, Cambridge, MA, USA*; YAN ZHOU, *JILA, National Institute of Standards and Technology and Univ. of Colorado Department of Physics, University of Colorado, Boulder, Boulder, CO, USA*; TIMOTHY J BARNUM, ROBERT W FIELD, *Department of Chemistry, MIT, Cambridge, MA, USA*.

Free induction decay detected chirped pulse millimeter wave spectroscopy of Rydberg-Rydberg transitions in atoms and molecules is a powerful and flexible method for characterizing the electronic structure of Rydberg states and determining the structure and dynamics of the ion-core. Complicating the use of this technique are the difficulties in reliably and repeatedly accessing not just the most information rich core-nonpenetrating states, but also the low- $\ell$  core-penetrating Rydberg states in the area of principal quantum number  $n^* > 35$ . Small transition moments and narrow linewidths for transitions between valence electronic states and high Rydberg states are the primary limiting factor. We demonstrate a simple method to avoid the problem entirely by using chirped pulse technology operating in the frequency range of 260-295 GHz, which allows us to sample a lower range of  $n^*$  values than before with comparable frequency resolution and accuracy as our previous W-band experiments. Further improvements to our experiment in order to accurately capture details of Stark demolition, a technique that provides rapid differentiation between core-penetrating and core-nonpenetrating states, will also be discussed.